

# Chronic Liver Disease and Cirrhosis Deaths

## Chronic Liver Disease and Cirrhosis Deaths

- Most recent data (2014): 9.1 deaths per 100,000, including 204 deaths.
- Trend over time: 2000-14: rising
- Description: Includes deaths from alcoholic liver disease, chronic hepatitis, and cirrhosis of liver
- Healthy People 2020 Goal: None listed for this indicator.

Chronic liver disease and cirrhosis deaths, King County, 2010-2014 average

	Rate	Lower CI	Upper CI	Count per Year
<b>King County</b> (all ages)	9.3	8.8	9.9	204
<b>AGE</b>				
<18	0.0	0.0	0.2	0
18-24	*	*	*	*
25-44	3.5	2.9	4.3	22
45-64	24.1	22.3	26.0	128
65+	23.8	21.1	26.8	54
<b>GENDER</b>				
Male	12.4	11.4	13.4	130
Female	6.6	5.9	7.3	74
<b>RACE/ETHNICITY</b>				
AIAN	48.8	34.5	69.8	9
Asian	2.8	2.0	3.9	8
Black	7.4	5.3	10.3	9
Hispanic	9.3	6.6	13.4	10
Multiple	3.9 §	1.7 §	9.8 §	2
NHPI	*	*	*	*
White	10.3	9.6	11.0	174
<b>NEIGHBORHOOD POVERTY</b>				
High	16.1	14.0	18.5	42
Medium	9.3	8.5	10	127
Low	5.8	4.9	6.8	31
<b>REGION</b>				
East	6.2	5.3	7.3	36
North	9.9	7.7	12.6	15
Seattle	10.1	9.0	11.3	67
South	10.5	9.5	11.6	82
<b>STATE</b>				
Washington State	11.0	10.7	11.4	853

### Comparisons:

- Higher than King County rate (yellow)
- Lower than King County rate (green)

### Notes:

Source: WAState DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.

\*Rate = cases per 100,000 population, age-adjusted to the 2000 US population.


CI is 95% Confidence Interval.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

## Chronic liver disease and cirrhosis deaths King County, 2010-2014 average



Source: WA State DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.

^Rate = cases per 100,000 population, age-adjusted to the 2000 US population.

|-----| Confidence interval shows range that includes true value 95% of the time.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

Poverty = Neighborhood poverty levels defined by median household income.

# Chronic liver disease and cirrhosis deaths by health reporting areas, King County, 2010-2014 average

	Rate	Lower CI	Upper CI	Count per Year
<b>King County (all ages)</b>	9.3	8.8	9.9	204
Auburn	16.7	12.6	21.9	12
..Auburn-North	15.7	10.7	22.3	7
..Auburn-South	18.3	11.7	28.0	5
Bear Creek/Carnation/Duvall	4.4	2.2	8.5	3
Bellevue	6.9	5.2	9.2	11
..Bellevue-Central	11.2	6.8	17.7	4
..Bellevue-NE	6.3	3.3	11.4	3
..Bellevue-South	4.5 §	2.0 §	10 §	2
..Bellevue-West	6.1 §	3.0 §	11.8 §	2
Black Diamond/Enumclaw/SE County	8.9	5.7	13.6	5
Bothell/Woodinville	9.2	5.1	15.5	3
Burien	14.6	10.5	20	8
Covington/Maple Valley	4.3	2.4	8.1	3
Des Moines/Normandy Park	12.5	8.2	18.6	6
East Federal Way	8.2	4.6	14.3	3
Fairwood	9.6	4.6	19.0	2
Federal Way	10.5	7.8	14.1	10
..Fed Way-Central/Military Rd	9.1	6.0	13.6	6
..Fed Way-Dash Point/Woodmont	13.2	8.1	20.8	4
Issaquah	5.7 §	2.7 §	11.5 §	2
Kenmore/LFP	6.9	3.8	12.0	3
Kent	9.0	6.7	11.9	11
..Kent-East	8.8	4.9	15.2	3
..Kent-SE	8.5	5.3	13.2	4
..Kent-West	10.1	5.6	18.7	3
Kirkland	7.4	5.1	10.5	7
..Kirkland	6.6	3.9	10.8	4
..Kirkland North	8.4	4.7	14.6	3
Mercer Isle/Pt Cities	4.5 §	1.5 §	11.7 §	1
Newcastle/Four Creeks	13.8	8.4	22.1	5
North Highline	17.1	9.9	29.0	3
Redmond	4.7	2.4	8.7	2
Renton	9.2	6.9	12.1	11
..Renton-East	4.3 §	1.7 §	10.2 §	1
..Renton-North	11.9	7.1	19.1	4
..Renton-South	11.0	7.3	16.0	6
Sammamish	3.9 §	1.3 §	10.1 §	2
SeaTac/Tukwila	13.6	9.2	19.9	6
Seattle	10.1	9.0	11.3	67
..Ballard	11.2	7.5	16.5	6
..Beacon/Gtown/S.Park	10.1	6.3	15.4	5
..Capitol Hill/E.lake	7.3	4.1	13.5	3
..Central Seattle	5.1 §	2.5 §	9.7 §	2
..Delridge	11.8	7.0	19.8	4
..Downtown	31.9	25.2	46.3	16
..Fremont/Greenlake	5.8	3.1	10.9	3
..NE Seattle	8.3	5.3	12.6	5
..North Seattle	7.3	4.2	12.2	4
..NW Seattle	10.4	6.7	16.2	5
..QA/Magnolia	4.4	2.5	8.1	3
..SE Seattle	8.9	5.3	14.2	4
..West Seattle	11.1	7.8	16.0	8
Shoreline	12.1	8.6	16.8	9
Snoqualmie/North Bend/Skykomish	7.4	3.8	14.1	3
Vashon Island	8.0 §	3.3 §	24.2 §	2

## Comparisons:

Higher than King County rate (yellow)

Lower than King County rate (green)

## Notes:

Source: WA State DOH, Center for Health Statistics, Death Certificates.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.

\*Rate = cases per 100,000 population, age-adjusted to the 2000 US population.


CI is 95% Confidence Interval.

\* Too few cases to protect confidentiality and/or report reliable rates.


§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

## Chronic liver disease and cirrhosis deaths by health reporting areas King County, 2010-2014 average



## Chronic liver disease and cirrhosis deaths King County, 2000-2014, three-year rolling averages



<sup>a</sup>Rate = cases per 100,000 population, age-adjusted to the 2000 US population.  
Trends over time: King County (2000-14: rising); East KC (2000-14: flat); North KC (2000-14: flat); Seattle (2000-14: flat); South KC (2000-14: rising); Washington State (2000-05: flat; 2005-14: rising)  
Source: WA State DOH, Center for Health Statistics, Death Certificates.  
Prepared by Public Health - Seattle & King County, APDE, 08/2016.

Produced by the  
Assessment, Policy Development & Evaluation Unit  
Public Health - Seattle & King County  
08/16